

Temporal Integration of Motion Streaks in Migraine

Louise O'Hare ^{1,†} 

¹ University of Lincoln; LOHare@lincoln.ac.uk

† These authors contributed equally to this work.

Version July 11, 2018 submitted to Vision

Abstract: Migraine is associated with differences in visual perception, specifically, deficits in the perception of motion. Migraine groups commonly show poorer performance (higher thresholds) on global motion tasks compared to control groups. Successful performance on a global motion task depends on several factors, including integrating signals over time. A “motion streak” task was used to investigate specifically integration over time in migraine and control groups. The motion streak effect depends on the integration of a moving point over time to create the illusion of a line, or “streak”. There was evidence of a slower optimum speed for eliciting the motion streak effect in migraine compared to control groups, suggesting temporal integration is different in migraine. In addition, performance on the motion streak task showed a relationship with headache frequency.

Keywords: migraine; temporal integration; motion perception

1. Introduction

Migraine is a common, debilitating neurological disorder affecting around 13% of the adult population [1]. The International Headache Society classification criteria [2] for migraine requires a minimum of five episodes of headache attacks that last 4–72 hours when untreated, with the following qualities: headache is located on one side of the head, the headache has a pulsating quality, moderate to severe pain, the headache is aggravated by physical activity. Additionally the headache is accompanied by either nausea/vomiting, and/or by aversion to light and sound (photophobia and phonophobia, respectively). Approximately 39% of those with migraine experience migraine aura, a set of sensory disturbances preceding the onset of the headache itself [3], and the most commonly reported modality for aura is visual [1]. There is a distinction between migraine with aura (MA) and migraine without aura (MO).

Migraine has strong associations with vision: around 80% of those with migraine are light hypersensitive [4]. Those with migraine tend to report extreme discomfort on viewing striped patterns in between migraine attacks (interictally) [5]. Additionally, there are many differences in performance between interictal migraine and control groups on a variety of visual tasks (see [6], for a review). One of the most robust findings is poorer performance of interictal migraine groups compared to control groups on global motion tasks (e.g. [7] [8] [9] [10] [11] [12] [13]).

Global motion tasks typically involve detecting the direction of a coherently moving field of dots (or other elements) against a background of noise elements, for example dots moving in random directions. Success on a global motion task depends on multiple processes: detecting the local motion signals, then integrating the motion signals, and finally discriminating the direction of the signal from background noise [14]. The ability to detect coherent motion in the absence of stimulus noise (i.e. detecting local motion) is unimpaired in migraine compared to control groups [13]. Therefore, it appears that either integration of motion and/or the ability to discriminate signal from background noise is impaired in those with migraine.

The motion streak effect depends on temporal integration: this is a visual phenomenon where a fast-moving point stimulus can be integrated over time to produce the appearance of a line in the direction of motion [15]. Streaks are also called 'speedlines' and are used in artworks to imply motion in the static image [16]. The visual system can exploit this streak effect to enable the use of orientation detectors to facilitate the detection of the direction of motion ([15][17] [18] [19] [20]). In short, a point light source moving rapidly is integrated to enable the use of orientation detectors to facilitate direction discrimination.

Orientation detectors are subject to iso-feature suppression, a form of inhibition from neighbouring detectors of a similar type [21], which are sharpened by short-range inhibitory processes [22]. Therefore, the ability to use orientation detectors to discern motion direction can be estimated by comparing the masking effects of a parallel vs orthogonal noise mask background, see Figure 1 for a schematic of the stimulus. Backgrounds oriented parallel to the motion direction will mask the streak effect, resulting in poorer performance ([15] [16][17], [18] [19]). Additionally, backgrounds orthogonal to motion direction can facilitate the motion direction discrimination [15].




Figure 1. Figure showing schematic diagram of the stimulus. The Gaussian dot is presented in the centre of the screen, and moves either up, down, left or right. The task is to identify dot motion and indicate this using the corresponding arrow keys. There were five levels of dot contrast (relative to midgrey) and five possible speeds (1, 4, 7, 10, 13 °/s) . The horizontal background orientation is shown, there is also a vertical background and homogenous grey background. Responses for motion orthogonal to the background were pooled, as were responses to motion parallel to the background. These were normalised against performance against the homogenous grey background.

In order to investigate temporal integration processes, the speed of the moving element was varied, as the most effective speed for the motion streak effect is dependent on the size and contrast of the moving element. The strength of the motion streak depends on the speed of the moving element

[15], the size of the moving element [15], the distance travelled by the moving element in a given display time (number of frames) [23]. A stronger motion streak effect leads to improved ability to discriminate motion direction ([15] [23]), and increasing the speed (up to a point) is one method of increasing the strength of the motion streak effect. There is individual variation in the critical speed for the motion streak effect (the speed at which the motion streak effect begins to appear), which will depend on the length of the temporal integration window for an observer.

Vertical as well as horizontal motion were estimated in the current study, as there are differences in optokinetic reflex eye movements in those with MA as well as those with migraineous vertigo, compared to control groups - those with MA show increased velocity for ocular following movements (the eyes move faster) compared to control participants [24]. Additionally, optokinetic stimulation has been demonstrated to increase motion sickness in those with migraine more than control participants [25]. The optokinetic effect is greater for horizontal compared to vertical motion in both non-human primates [26], and in humans [27]. As there is a difference in the strength of the optokinetic effect in the horizontal compared to vertical directions, and this is different in migraine and control groups, it is possible that any results could be accounted for by eye movements. To control for this possibility, it was important to estimate the effect in both planes (including all four directions of up, down, left and right).

To summarise, the motion streak effect relies specifically on temporal integration processes. Motion streak in the presence of a background oriented parallel to the motion direction will result in the greatest threshold elevation (poorest performance). This elevation in thresholds is thought to be due to orientation-specific inhibitory processes that arise due to the integration of the motion signal into a line [15]. It is expected that there will be a difference between migraine and control groups for the motion streak task, if there are differences in temporal integration processes in those with migraine. Further, it is expected that there will be differences in the optimum speed for the motion streak effect between the migraine and control groups, because the optimum speed depends on the relative speed of the motion compared to the temporal integration window.

2. Results

2.1. Orthogonal motion

Figure 2 A shows threshold elevation for five speeds of dot motion (1, 4, 7, 10 and 13°/s) against orthogonal noise backgrounds (elevation compared to homogenous grey background). Mixed ANOVA was conducted with group as a between subjects factor, and speed as a within-subjects factor, degrees of freedom were adjusted where the assumption of homogeneity of variance was not met (see Materials and Methods for more details). There was a significant main effect of speed ($F(2.95, 144.57) = 4.23$, $p = 0.007$, $\eta_G^2 = 0.06$). Post-hoc least-squares means contrast were calculated using the package "lsmeans" ([28], for details see Methods section), using the Bonferroni correction for ten comparisons. There was a significant difference between 4 and 13°/s ($p = 0.0040$), none of the other comparisons were statistically significant after correction for multiple comparisons. Threshold elevation was overall greater for the migraine group compared to the control group ($F(1, 49) = 5.70$, $p = 0.02$, $\eta_G^2 = 0.03$). There was no statistically significant interaction between group and speed ($F(2.95, 144.57) = 2.05$, $p = 0.11$, $\eta_G^2 = 0.03$).




Figure 2. Figure showing threshold elevation (dB) against speed of dot motion for migraine and control groups. Error bars are one standard error of the mean: (a) Threshold (dB) against speed (deg/s) for motion orthogonal to the background orientation. (b) Threshold (dB) against speed (deg/s) for motion parallel to the background orientation.

2.2. Parallel motion

Figure 2B shows threshold elevation (dB) for motion against parallel filtered noise backgrounds (elevation was calculated by dividing threshold against filtered noise background by threshold for homogenous grey background). Mixed ANOVA was conducted with group as a between subjects factor, and speed as a within-subjects factor (see Materials and Methods for more details). There is a significant main effect of speed ($F(3.08,150.86) = 6.33, p = 0.0004, \eta_G^2 = 0.09$). There is no effect of group ($F(1,49) = 0.07, p < 0.80, \eta_G^2 = 0.0003$). There is an interaction between migraine and speed ($F(3.08,150.86) = 4.06, p = 0.008, \eta_G^2 = 0.06$). There was a significant difference between migraine and control groups at $10^\circ/\text{s}$ ($p = 0.044$) and at $13^\circ/\text{s}$ ($p = 0.011$), none of the other comparisons were statistically significant when adjusting for multiple comparisons using Bonferroni correction.

2.3. Difference in threshold elevation

Figure 3 shows the difference in threshold elevation between orthogonal and parallel backgrounds, to highlight the cost of the parallel background compared to the orthogonal background. There is a significant main effect of speed ($F(2.89,141.78) = 34.04, p < 0.0001, \eta_G^2 = 0.34$). Post-hoc comparisons showed a significant difference between 1 and $7^\circ/\text{s}$ ($p = 0.0093$), between 1 and $10^\circ/\text{s}$ ($p < 0.0001$), and between 1 and $13^\circ/\text{s}$ ($p < 0.0001$). There was also a significant difference between 4 and $10^\circ/\text{s}$ ($p < 0.0001$), and between 4 and $13^\circ/\text{s}$ ($p < 0.0001$). There was a significant difference between 7 and $10^\circ/\text{s}$ ($p = 0.0005$), and between 7 and $13^\circ/\text{s}$ ($p < 0.0001$). The other comparisons were not statistically significant when adjusting for multiple comparisons using the Bonferroni correction. This indicates

that the increase in masking between orthogonal and parallel backgrounds depends on speed. The difference between orthogonal and parallel backgrounds is smaller for the migraine group compared to the control group ($F(1,49) = 5.40$, $p = 0.02$, $\eta_G^2 = 0.03$). The interaction between speed and migraine group is not statistically significant ($F(2.89,141.78) = 2.63$, $p = 0.05$, $\eta_G^2 = 0.04$), however the effect size estimate indicates a small effect.




Figure 3. Figure showing difference in threshold elevation (parallel-orthogonal background) against speed for migraine and control groups. Error bars are one standard error of the mean.

2.4. Relationship to headache frequency

Frequency of headaches was assessed using a questionnaire. Participants responded by choosing one of the following options - whether their headaches occurred never, less than once a month, 1 to 3 times a month, 3 to 10 times a month, or more than 10 times a month. Estimated frequency of headaches was coded to take the midpoint of each category of answers, for example, 1-3 times a month was recoded as 2, the midpoint. Values less than 1 were coded as 0.5, values of 10 or more were coded as 10. A linear mixed effects model was fitted to see if headache frequency of the migraine group predicted the difference in threshold elevation (orthogonal minus parallel, as in 3). The linear mixed effect model has some advantages in terms of non-independent observations, for a discussion, see [29]. An alternative model included headache frequency and dot speed as fixed factors, and also observer as a random factor. The null model included only the factors of dot speed as a fixed factor, and observer as a random factor. Thus, two models were compared - the alternative model includes headache frequency, and the null model does not include headache frequency as a predictor. The ability to predict the difference in threshold elevation was significantly better for the alternative model (with headache frequency) compared to the null model (without headache frequency) ($\chi^2(1) = 8.01$, $p = 0.0046$). The difference in threshold elevation motion against orthogonal and parallel backgrounds

increased with headache frequency by 0.32dB (standard error = 0.11). This indicates a greater threshold elevation with those who experience more frequent headaches.

3. Discussion

The motion streak task specifically was chosen as this effect depends on temporal integration processes [15]. Importantly, backgrounds orthogonal to motion direction should present less masking than backgrounds parallel to motion direction [15]. For both groups, the experiment partially replicated the motion streak effect originally found by [15] - a detrimental effect of the background oriented parallel to the motion direction, which can be seen in Figure 2B.

Importantly, there are group differences shown on the motion streak task; those with migraine showed poorer performance for orthogonal backgrounds. For motion against parallel backgrounds, there is an interaction between migraine and speed, with a bigger motion streak effect in the control group compared to the migraine group at faster speeds. From Figure 2 B it can be seen that the migraine group reach asymptote at slower speeds compared to the control group. This suggests the optimum speed for the motion streak is slower for the migraine compared to the control group. To highlight the motion streak effect, threshold elevation against orthogonal and parallel backgrounds was compared, a greater difference would be evidence of a greater motion streak effect, which can be seen in Figure 3. Figure 3 shows a lower difference between motion parallel and orthogonal to the background for the migraine group compared to the control group, indicating a weaker motion streak effect. Therefore it may be the case that there is a shorter temporal integration window in migraine, as the streak effect occurs earlier, and is weaker, in this group.

The motion streak effect (difference in threshold elevation for motion against orthogonal backgrounds compared to motion against parallel backgrounds) was estimated in both vertical and horizontal motion direction. Both motion orientations were used to control for the possibility of anisotropy in performance, as there is evidence that the optokinetic effect is different in migraine compared to control groups [24] [25]. The motion streak effect was found in both horizontal and vertical motion directions (see Appendix B), and so both directions of motion (set against orthogonal and parallel backgrounds) were used in the main study. When direction of motion (horizontal and vertical dot motion) was included as a factor in the analysis of the difference in threshold elevation between orthogonal and parallel backgrounds, there was some evidence of poorer performance overall for the vertical motion direction compared to the horizontal motion direction, but no group-specific effect of direction of motion. Therefore, the poorer performance for vertical compared to horizontal dot motion may be evidence of the optokinetic effect, but this appears to be no different in the migraine and control groups in the current study.

3.1. Relationship to migraine characteristics

Headache frequency (in the migraine group) was predictive of a greater difference in threshold elevation for motion against the orthogonal compared to the parallel backgrounds. Those with more frequent headaches tend to show a bigger difference, and so bigger motion streak effects. A relationship between attack frequency and cognitive test performance, including tasks of visual memory has been shown previously e.g. [30]. Additionally, those with migraine show elevated response amplitude when measured over the somatosensory cortex, which is positively correlated to the frequency of migraine attacks [31]. However, the relationship between headache frequency and cognitive test performance is mixed (see [32] for a review). Visual field deficits were not found to correlate with attack frequency [33] [34].

The current study only considers interictal migraine - those who had experienced a migraine attack too recently were excluded from the study. Effects of the migraine cycle would be interesting to investigate in future research, as there are effects of the migraine cycle shown in both behavioural e.g. [35] [36] and electrophysiological [37] [38] [39] responses. Due to multiple testing sessions it was not possible to estimate effects of the migraine cycle in the current experiment. The use of a staircase

procedure would enable thresholds to be obtained more rapidly (therefore no longer needing multiple testing sessions) and also avoid potential problems with floor effects.

3.2. Temporal integration mechanisms

The integration of two (or more) rapidly presented stimuli is of particular importance in migraine, as those with migraine report aversion on viewing flickering stimuli e.g. [40] [41]. It has been suggested that flickering stimuli, such as the "two flash" test is a simple estimate of temporal integration, without the spatial component that is inherent in motion perception [20]. The critical flicker fusion threshold is the lowest frequency at which flickering light will be seen as detectable, rather than a steady light source. Previous studies reported lower flicker fusion thresholds in migraine compared to control groups [42] [43], which would be indicative of increased summation of responses over time. In the case of the [42] paper, there is a significant difference in the flicker fusion threshold between MO and control groups, but not between MA and control groups. This is not in agreement with findings that those with migraine are better able to detect briefly presented stimuli compared to control group [44], or the current study showing a weaker motion streak effect in the migraine group. It is unclear why this might be the case. Speculatively, it has been shown that habituation to repetitive stimuli is weaker in migraine compared to control group, indicating that there is a sustained response to repetitive stimuli in the migraine group [45]. This could be an important difference between temporal integration of single stimuli compared to repetitive stimulation.

Brain connectivity is important for the integration of sensory information, and this has been suggested to be a potential cause of sensory processing differences seen in those with migraine [46]. Local functional connectivity within visual areas is unimpaired in those with migraine [46]. Further, connectivity in the resting state has been shown to be increased in those with MA compared to control groups [47]. Temporal integration of multisensory stimuli has been linked to functional connectivity in EEG studies [48] [49]. Functional connectivity in terms of coherent neural oscillations has been shown to be different in those with migraine compared to control groups [50]; those with MA showed stronger beta band coherence compared to MO and control groups, whereas those with MO showed weaker alpha band (8-12Hz) coherence compared to MA and control group. Alpha band (8-12Hz) oscillations in the occipital areas of the brain are thought to control the timing of information processing in the visual areas of the brain [51]. The alpha band oscillations provide a "window of excitability" [52]; if alpha power is high at stimulus onset, then it is unlikely to be perceived, conversely, if occipital alpha power is low at stimulus onset then it is more likely to be perceived [53] [54] [55]. Increasing alpha band frequency has been demonstrated to have effects on the temporal integration window in normal observers for multisensory integration studies [56] [57]. Evidence for a longer integration window has been demonstrated in multimodal integration between the visual and vestibular system in those with migraine [58]. The current study shows some evidence that the temporal integration window is different in migraine compared to control groups, and this task is thought to depend on integration of the signal over time in order for the streak to appear. In particular, there is evidence of a slower optimum speed for the migraine group compared to the control group, and so it might be thought that the neural oscillations may also be slower in those with migraine compared to control groups. In future research, it would be useful to assess whether temporal integration relates to coherence of neural oscillations (functional connectivity) in those with migraine, as this has clear predictions for clinical outcomes and potential treatments [59].

3.3. Limitations of the current study

In the current experiment, any potential group differences in motion detection, or lateral inhibition *per se* were controlled for by considering threshold elevation between oriented backgrounds and homogenous grey backgrounds. Previous research has shown increased lateral inhibition in migraine compared to control group using behavioural measures [60]. Additionally, electrophysiological data has also shown differences in short-range lateral inhibition in those with migraine [61]. However, there

is evidence [44] [62] that the short-range lateral interactions are no different in migraine compared to control groups. Importantly, if there were a difference in short-range interactions, both vertical and horizontal background conditions should be equally affected, which was not the case. Additionally, if the migraine group is simply more susceptible to noise from the addition of an oriented background compared to control group, then this should affect both backgrounds (orthogonal and parallel to the direction of motion) equally, and so no specific motion streak effect should be seen. Again, this was not the case. It is also possible that differences in the ability to detect motion in migraine - for example, differences in the opponent motion system [63]. There is increased cortical thickness in area V3A and area MT+, both of which are associated with motion processing [64], which might be expected to result in potential differences in motion processing in migraine compared to control groups. However, there is evidence to suggest that the detection of motion in the absence of noise is unimpaired in migraine groups [13]. Even so, the difference in threshold elevation between the parallel and orthogonal the background was estimated to limit the effect of differences in motion detection between the two groups accounting for the results in the current experiment.

One limitation of the current study is that not all individuals had a diagnosis of migraine by a medical professional. It has been suggested that many of those with migraine do not seek medical advice, estimates vary between 37% [65] and around half [66]. Participants in the current study were asked to complete a questionnaire to assess whether they fulfilled the International Headache Society criteria [2], diagnosis was not made by a trained neurologist. In order to address this concern, Appendix F contains a second analysis of the results, including only those individuals with a diagnosis of migraine from a medical professional. There is a similar pattern of results when only these individuals are included in the migraine group. A review of cognitive deficits in migraine showed that migraine with aura was inconsistently associated with cognitive deficits, including performance on visually-based tasks, in migraine [32]. Unfortunately, in this study there were insufficient participants to be able to split the migraine group into MA and MO groups. This is a limitation of the study as it is not clear if these two groups differ in terms of their performance on this task. Those with MA do show differences to those with MO e.g. [50], but this is not always the case, for example, other studies have not shown differences in motion perception between MA and MO groups e.g. [67] [8] [9] [68] [13].

Another limitation is the reliance on self-report for the measures of normal visual acuity. It is possible that individuals have uncorrected visual deficits that they may or not be aware of. It has been shown previously that differences in contrast sensitivity also can correlate with performance on motion based tasks in those with migraine e.g. [12]. Poorer visual acuity and/or differences in contrast sensitivity could account for the overall poorer performance in the migraine group compared to the control group. However, the difference in threshold elevation cannot be accounted for by poorer acuity or reduced contrast sensitivity, as either of these would result in the same decrease in both conditions, not an orientation-specific effect of the background.

Recent fMRI studies demonstrate that brain networks associated with attention have been shown to be different in those with migraine compared to control group in between migraine attacks e.g. [69]. Additionally, [70] showed reduced thalamocortical connectivity in MO compared to control group at resting state during a migraine attack. Other researchers have also demonstrated reduced activity at rest in networks associated with attention in MO, but not MA, compared to control groups [71]. If attention is different in those with migraine compared to control group, then this might be expected to account for differences in susceptibility to externally added noise. Differences in attentional network connectivity could account for the overall poorer performance in the migraine group compared to the control group in the current study. However, differences in attention alone cannot account for the difference in threshold elevation between the orthogonal and parallel backgrounds.

4. Materials and Methods

4.1. Observers

60 observers with corrected-to-normal vision were recruited to the study using advertising around the University campus. Visual acuity was based on self-report, if participants reported needing optical corrections, they were encouraged to wear them. All experiments adhered to the World Medical Association Declaration of Helsinki (2013) and were scrutinised by the University of Lincoln School of Psychology Ethics committee, ethical approval code PSY1617240. Written informed consent was obtained from all observers before commencing the study.

Migraine was defined as fulfilling the International Headache Society diagnostic criteria [2]. This was determined via a questionnaire, administered by the experimenter. Not all individuals had a migraine diagnosis from a medical professional (e.g. GP or neurologist), but many did. The diagnosis by a medical professional can be seen in Table 1. Individuals with 'probable migraine', who experience an insufficient absolute number of headache attacks (fewer than 5 attacks) were excluded. Five observers did not fulfill the criteria. Two of these five who were excluded as they did not fulfill the IHS criteria had a diagnosis of migraine by a medical professional. These were excluded from the analysis here, but included in the analysis of only those with a professional diagnosis of migraine in the Appendix. Two additional observers were excluded as they had a diagnosis of tension headache, and chronic headache, respectively. Those with a headache attack within 2 days were excluded (one observer). Control groups contained individuals who had reported no migraine, and no regular headaches. One control participant was excluded due to experiencing too many headaches. As the migraine classification was based on self-report, the experimenter would have known which participants were in the migraine and which were in the control group. In order to minimise subjective bias from this experimenter knowledge, all participants completed all conditions of the experiments, and all trials were presented in random order, by computer, therefore minimising subjective bias. After exclusions there were 29 control participants (mean age 23.517 years, SD = 6.399), eight of whom were male, and 22 migraine participants (mean age 24.82 years, SD = 8.16), three of whom were male.

Table 1. Table showing characteristics of those individuals with migraine included in the final analysis, after exclusions. Monthly freq = monthly frequency of headaches, Visual = are there any visual disturbances during the attack, Speech = are there any speech disturbances during the attack, Prof diagnosis = do you have a diagnosis of migraine, migraine with aura, migraine without aura by a medical professional (either GP or neurologist), M = migraine, MO = migraine without aura, MA = migraine with aura, AB = abdominal migraine.

Gender	Age	Monthly freq	Visual	Speech	Prof diagnosis	Last attack
F	28	<1	yes	yes	MA	2 weeks
M	41	< 1	yes	no	no	2 months
F	23	1 to 3	yes	no	M	8 days
F	48	<1	yes	no	MA	18 months
M	19	3 to 10	yes	no	MA	2 weeks
F	29	<1	yes	yes	MA	1 month
F	18	<1	yes	no	MA	4 months
F	22	1 to 3	no	yes	no	2 weeks
F	19	3 to 10	yes	no	M	5-6 days
F	21	1 to 3	yes	no	MO	2-3 weeks
F	21	10+	yes	yes	MA	3-4 days
F	22	<1	yes	yes	AB	months
F	19	1 to 3	no	no	M	6 days
F	20	1 to 3	yes	no	M	1 week
F	21	3 to 10	yes	no	no	1 week
F	18	1 to 3	yes	no	MA	2 weeks
F	22	1 to 3	yes	no	M	6 days
F	23	1 to 3	yes	no	no	3 weeks
F	21	3 to 10	yes	no	MO	2 days
F	26	1 to 3	yes	no	M	2 weeks
M	24	<1	yes	no	no	2 months
F	41	<1	no	no	M	2 weeks

4.2. Apparatus

All experiments took place in a sound-attenuated, darkened room. An MSI (MS-7788) computer with i7-3990CPU Intel processor, NVida GeForce GTX 650 graphics card, and a 64-bit Windows 7 operating system was used to create and display stimuli. Stimuli were displayed using a 22-inch Illyama HM204DTA Vision Master Pro 514 Diamondtron U3-CRT monitor, calibrated with LS100 Minolta photometer. A Bits Sharp signal processor (Cambridge Research Systems, Cambridge, UK) was used to convert the RGB signal into 14-bit greyscale. Mean luminance of the display used for the behavioural task was 44.24 cd/m². Refresh rate was 85Hz, resolution was 1024 x 768 pixels. Viewing distance was 100cm, head movements were restricted using a chinrest. Matlab Version 2015a (The Mathworks, Natick, USA) and the Psychtoolbox ([72] [73] [74]) were used to generate and display stimuli.

4.3. Stimuli

The stimulus was a Gaussian blob (0.1°, or 6 arcmin, of visual angle, measured across two standard deviations). Dot speed was 1, 4, 7, 10, 13°/s. The dot always began in the centre of the screen. The dot could move either horizontally, or vertically. There were again five levels of dot contrast, between 0% and 10% contrast, compared to the homogenous mid-grey background. There were three background types, no background (homogeneous grey), filtered noise oriented parallel to the motion, and filtered noise oriented perpendicular to the motion. The filter was a raised radial cosine function, see equation 1.

$$H(f) = \begin{cases} T \\ \frac{T}{2} [1 + \cos(\frac{\pi T}{\beta})(|\log(f) - \log(f_0)| - \frac{1-\beta}{2T})] \\ 0 \end{cases}$$

$$\text{for } \begin{cases} (0 \leq |\log(f) - \log(f_0)| \leq \frac{1-\beta}{2T}) \\ (\frac{1-\beta}{2T} \leq |\log(f) - \log(f_0)| \leq \frac{1+\beta}{2T}) \\ (|\log(f) - \log(f_0)| > \frac{1+\beta}{2T}) \end{cases} \quad (1)$$

where T is 0.9, β is 0.5, and f_0 is the centre frequency is of 2.9 cycles/°. The bandwidth of the filtered noise was relative to the dot size, the centre frequency $\pm 0.1^\circ$ from the centre frequency. Stimuli were presented in a Gaussian-edged window with an aperture of 5.06° and a soft edge $\sigma = 0.51^\circ$. A schematic stimulus can be seen in Figure 1. There were 3 backgrounds \times 5 contrast levels \times 5 speeds \times 20 repetitions of each trial resulting in a total of 1,500 trials. Trial order was randomised for each participant individually. Observers indicated the direction of motion in a four-alternative forced-choice procedure, choosing either left, right, up or down on any given trial, by pressing the corresponding arrow keys. The experiment was divided into five blocks (300 trials in each block), over three sessions. Each session lasted around 1 hour. The first block was used as a practice session and excluded from the analysis.

Stimulus parameters were chosen based on pilot work, the first pilot study was to estimate the dot size and contrast levels needed for the main experiment, using two dot sizes and horizontal motion only. Results from this can be seen in Appendix A. The second pilot study was to ensure that the stimulus parameters from the first pilot study were suitable for both horizontal and vertical motion directions. Results from this can be seen in Appendix B.

4.4. Statistical analysis

The level of dot contrast needed to achieve 75% correct performance was estimated for each of the conditions. Threshold performance (against filtered noise background) was normalised to the baseline (homogenous grey background) performance, in order to assess the effect of motion streaks, and not just performance against a mask. Normalisation was achieved by dividing the oriented background (horizontal or vertical) by the homogenous grey background. Normalised thresholds are expressed in dB. This was achieved using the following equation:

$$20\log_{10}(\text{thresholdOriented} / \text{thresholdHomogenousGrey}) \quad (2)$$

Where thresholdOriented is the threshold obtained for motion against the oriented background, and $\text{thresholdHomogenousGrey}$ is the threshold obtained against the homogenous grey background. Inferential statistics were calculated using R [75] with packages “afex”, [76], “lsmeans” [28], and “lme4” [77]. Mixed ANOVA was conducted with migraine as a between subjects factor, and background and speed as within-subjects factors. Data were not normally distributed according to a Shapiro-Wilk test: for motion against orthogonal backgrounds ($W=0.90$, $p = 6.86 \times 10^{-12}$), for motion against parallel backgrounds ($W=0.94$, $p = 2.16 \times 10^{-8}$), for difference in threshold elevation ($W = 0.87$, $p = 8.14 \times 10^{-14}$). However, the ANOVA can be robust to certain violations of the assumption of normality, depending on the severity of the violation [78], [79], [80]. Violations of these assumptions also applies to the post-hoc tests. For completeness, non-parametric versions of the tests were performed using the R package “nparLD” [81], using the Wald test statistic, and results showed that there were no differences in the overall pattern of results between the non-parametric versions of the analysis.

5. Conclusions

The motion streak task was used as an estimate of temporal integration in migraine and control groups. Those with migraine showed poorer performance overall, suggesting an inability to exclude external noise from the background in migraine. For motion against parallel backgrounds, there was a more pronounced effect of dot speed for the control group compared to the migraine group, and importantly evidence of motion streak effects at slower speeds for the migraine group. This suggests a difference in temporal integration in the migraine compared to the control group.

Acknowledgments: This work was supported by the University of Lincoln College Research Fund. Thanks to Andrea Pavan, Petra Pollux and Stefano Belli for their comments on the draft.

Author Contributions: LOH designed and conducted the experiments, analysed the data, and wrote the paper.

Conflicts of Interest: The author declares no conflict of interest.

Abbreviations

The following abbreviations are used in this manuscript:

IHS	International Headache Society
M	migraine
MA	migraine with aura
MO	migraine without aura
tACS	transcranial alternating current stimulation

Appendix A Experiment One

Experiment 1 was to determine the stimulus parameters for the main study. Previous work on the motion streak has been conducted with limited numbers of experienced psychophysical observers e.g. [15], and so their threshold performance may be different to the target group for this experiment. Experiment 1 consisted of only one motion speed, but two dot sizes, to test that the motion streak occurred in both those with migraine and control group, using these stimulus parameters.

Appendix A.1 Method

Appendix A.1.1 Observers

52 participants with corrected-to-normal vision (based on self-report only) took part in the study. Migraine was defined as fulfilling the IHS criteria (2013), or had a diagnosis from a medical professional. Individuals with 'probable migraine' (insufficient frequency of headache attacks (fewer than 5 attacks) were excluded. One observer was excluded due to being diagnosed with cluster headache, rather than migraine. Two observers were excluded due to previous head trauma. Five further observers were excluded as their most recent headache attack was within 2 days. A final 3 observers had outlying thresholds (defined as exceeding 2 standard deviations of the mean), and so were excluded. 15 migraine and 20 control participants were included in analysis, after exclusions, see table below for migraine characteristics. The migraine group had an average age of 21.20 years (standard deviation 5.00), two were male. The control group had an average age of 20.7 (standard deviation 2.45) years, 6 of them were male.

Table A1. Individuals with migraine included in the first pilot study, after exclusions. These individuals are not the same group of people as those who took part in the main experiment. Monthly freq = monthly frequency of headaches, Visual = are there any visual disturbances during the attack, Speech = are there any speech disturbances during the attack, Prof diagnosis = do you have a diagnosis of migraine from a medical professional (e.g. GP or neurologist), M = migraine, MO = migraine without aura, MA = migraine with aura.

Gender	Age	Monthly freq	Visual	Speech	Prof diagnosis	Last attack
F	19	1-3	no	no	M	1 month
F	18	10+	yes	no	MA	2 months
F	19	0	yes	no	M	1 week
F	19	1-3	yes	no	MA	1 week
F	20	1-3	yes	no	no	2 weeks
F	15	1-3	yes	no	M	1 week
F	18	< 1	yes	no	MA	3 months
M	19	3-10	yes	no	no	2 weeks
F	19	< 1	yes	no	no	2 days
F	20	0	yes	no	MA	2 weeks
F	18	3-10	no	no	no	2 months
F	32	1-3	no	no	MO	1 month
M	24	1-3	no	no	no	1 week
F	31	< 1	yes	no	MA	12 months
F	27	< 1	yes	yes	MA	6 weeks

Appendix A.1.2 Stimuli

There were two sizes of Gaussian dot used, 0.1° and 0.2° . Dot speed was $13.02^\circ/\text{s}$. There were five levels of contrast or the Gaussian blob, varying from 0 % to 2.5 % compared to midgrey. There were 600 trials in total, 300 trials for each dot size.

Appendix A.2 Results

Figure A1 shows threshold elevation for migraine and control groups for two dot sizes, 0.1° and 0.2° . Note the difference in scale. For the smaller dot size, an effect of background ($F(1,33) = 27.95$, $p < 0.0001$, $\eta_G^2 = 0.18$), but for the larger dot size there was no effect of background ($F(1,33) = 0.55$, $p = 0.46$, $\eta_G^2 = 0.005$). This is due to ceiling effects with the larger dot size, as threshold elevation was very low.




Figure A1. Figure showing threshold elevation for (left) 0.1° dot size and (right) 0.2° dot size for migraine and control groups against horizontal and vertical backgrounds. Dot motion is horizontal. Error bars are one standard error of the mean.

Appendix B Experiment Two

Experiment 2 manipulated the direction of dot motion, between horizontal and vertical. This experiment was to check that the motion streak effect seen in Experiment 1 could be replicated, that it is not specific to one direction, and that the 4AFC design is suitable. 4AFC has the advantage over 2AFC as the likelihood of guessing correctly on any individual trial is reduced. There have been reports of anisotropies in motion discrimination e.g. [82] [83], and so the inclusion of the vertical direction of motion is to see that the motion streak can also be elicited in this direction with the same parameters.

Appendix B.1 Method

Appendix B.1.1 Observers

42 participants with corrected-to-normal vision (based on self-report only) took part in the study, again migraine was defined as fulfilling the IHS (2013) classification criteria. One observer was excluded as they had cluster headache, rather than migraine. One observer was excluded as their migraines are the result of a diagnosed brain disorder (pernicious anemia). One observer was excluded due to reporting a headache within 2 days of the testing session. Four observers in the migraine group, and one in the control group, were excluded as thresholds could not be fitted to their data, due to floor effects. 10 migraine and 16 control participants were included in the final analysis, after exclusions (see table below for details). The average age of the migraine group was 19.60, years

(standard deviation 1.51), one member of the migraine group was male. There were 16 individuals in the control group, mean age was 20.75 (standard deviation 2.91), 6 of whom were male.

Table A2. Individuals with migraine who took part in the second pilot experiment, after exclusions. These individuals are not the same group of people who took part in the main experiment, or the first pilot experiment. Monthly freq = monthly frequency of headaches, Visual = are there any visual disturbances during the attack, Speech = are there any speech disturbances during the attack, Prof diagnosis = do you have a diagnosis of migraine by a medical professional (e.g. GP or neurologist), M = migraine, MO = migraine without aura, MA = migraine with aura.

Gender	Age	Monthly freq	Visual	Speech	Prof diagnosis	Last attack
F	21	3-10	yes	no	M	1 week
F	18	3-10	yes	no	M	2 days
F	19	1-3	no	no	M	5 days
F	19	3-10	yes	no	no	3 days
F	19	< 1	no	no	no	2 days
F	20	1-3	no	no	no	2 weeks
F	18	1-3	no	no	no	1 week
F	20	3-10	yes	no	M	1 day
F	19	1-3	yes	no	MA	1 month
M	23	1-3	yes	no	no	2 weeks

Appendix B.1.2 Stimuli

The task was 4AFC, the direction of motion could be up, down, left and right, observers indicated the direction of motion using the corresponding arrow keys. Contrast of the dot was increased slightly from Experiment 1, to 5 levels of contrast between 0 and 10 % compared to midgrey. There were again 600 trials in total, this time 300 trials for each direction of motion.

Appendix B.2 Results

Figure A2 shows threshold elevation for migraine and control groups for horizontal and vertical motion. There was an effect of background for both horizontal ($F(1,24) = 21.88$, $p < 0.0001$, $\eta_G^2 = 0.17$) and vertical backgrounds ($F(1,24) = 36.27$, $p < 0.0001$, $\eta_G^2 = 0.20$). The motion streak effect is not specific to horizontal motion and so both motion directions were used in the full study.




Figure A2. Figure showing threshold elevation for (left) horizontal motion and (right) vertical motion against vertical and horizontal backgrounds. Dot size is 0.1° . Error bars are one standard error of the mean.

Appendix C Diagnosis of Migraine by a Medical Professional Only

The analysis was repeated using only those individuals with a professional diagnosis of migraine. There were 20 individuals with a professional diagnosis of migraine, mean age was 23.90 years (standard deviation 7.76), two of this group were male. There were 29 control group, mean age 23.52 years (standard deviation 6.39), 8 of this group were male. It is important to note, that many of these are the same individuals as in the main experiment, for clarity, the individuals included in this re-analysis are listed in Table A3. There were some cases where participants were not included in the original analysis (as they did not meet the IHS criteria) but have a professional diagnosis of migraine.

Table A3. Individuals with a professional diagnosis of migraine only. These data are from the main experiment, this time excluding those individuals who did not have a diagnosis of migraine by a medical professional. Monthly freq = monthly frequency of headaches, Visual = are there any visual disturbances during the attack, Speech = are there any speech disturbances during the attack, Prof diagnosis = do you have a diagnosis of migraine, migraine with aura, migraine without aura by a medical professional (either GP or neurologist), M = migraine, MO = migraine without aura, MA = migraine with aura, VM = vestibular migraine, AB = abdominal migraine.

Gender	Age	Monthly freq	Visual	Speech	Prof diagnosis	Last attack
F	28	<1	yes	yes	MA	2 weeks
F	23	1 to 3	yes	no	M	8 days
F	48	<1	yes	no	MA	18 months
M	19	3 to 10	yes	no	MA	2 weeks
F	29	<1	yes	yes	MA	1 month
F	18	<1	yes	no	MA	4 months
F	19	0	yes	no	VM	8-10 days
F	19	3 to 10	yes	no	M	5-6 days
M	23	0	yes	yes	M	7-9 months
F	21	<1	yes	no	M	1 month
F	21	1 to 3	yes	no	MO	2-3 weeks
F	21	10+	yes	yes	MA	3-4 days
F	22	<1	yes	yes	AB	months
F	19	1 to 3	no	no	M	6 days
F	20	1 to 3	yes	no	M	1 week
F	18	1 to 3	yes	no	MA	2 weeks
F	22	1 to 3	yes	no	M	6 days
F	21	3 to 10	yes	no	MO	2 days
F	26	1 to 3	yes	no	M	2 weeks
F	41	<1	no	no	M	2 weeks

4.34 Appendix C.1 Threshold elevation for motion against orthogonal backgrounds

4.35 There was a significant main effect of group ($F(1,47) = 7.09$, $p = 0.01$, $\eta_G^2 = 0.03$). There was a
 4.36 significant main effect of speed ($F(2.95,138.53) = 3.96$, $p = 0.01$, $\eta_G^2 = 0.06$). Post-hoc tests with Bonferroni
 4.37 correction for multiple comparisons showed there to be a significant difference between 1 and 13°/s, (p
 4.38 = 0.042), between 4 and 13°/s ($p = 0.015$), none of the other comparisons were statistically significant.




Figure A3. Figure showing difference in threshold elevation (parallel-orthogonal background) against speed for migraine and control groups. Error bars are one standard error of the mean.

Appendix C.2 Threshold elevation for motion against parallel backgrounds

For motion against parallel backgrounds, there is a significant main effect of speed ($F(3.07, 144.38) = 5.43$, $p = 0.001$, $\eta_G^2 = 0.08$), and a significant interaction between migraine group and speed ($F(3.07, 144.38) = 4.46$, $p = 0.005$, $\eta_G^2 = 0.07$). Post-hoc tests showed there to be a significant difference between control and migraine groups at $1^\circ/\text{s}$ ($p = 0.030$), and at $13^\circ/\text{s}$ ($p = 0.0076$), none of the other comparisons were statistically significant when adjusted for multiple comparisons using the Bonferroni correction.

Appendix C.3 Difference in threshold elevation between orthogonal and parallel backgrounds

There was a significant main effect of group on the difference in threshold elevation between orthogonal and parallel motion ($F(1, 47) = 4.82$, $p = 0.03$, $\eta_G^2 = 0.03$). There was also a significant main effect of speed ($F(3.13, 147.02) = 33.63$, $p < 0.0001$, $\eta_G^2 = 0.34$). There was a significant interaction between migraine group and speed ($F(3.13, 147.02) = 4.06$, $p = 0.007$, $\eta_G^2 = 0.06$). There was a significant difference between the migraine and control group at $13^\circ/\text{s}$ ($p < 0.0001$), none of the other comparisons were statistically significant when corrected for multiple comparisons using the Bonferroni correction.




Figure A4. Figure showing difference in threshold elevation (parallel-orthogonal background) against speed for migraine and control groups. Error bars are one standard error of the mean.

Appendix C.4 Relationship to headache frequency

A general linear mixed effects model was created to predict the difference in threshold elevation between orthogonal and parallel backgrounds, including headache frequency and speed as a fixed factors, and subject as a random factor. This was compared to a null model that included only speed and subject as factors. There was a relationship between headache frequency and the difference in threshold elevation ($\chi^2(1) = 4.23$, $p = 0.040$). This increased the difference in threshold elevation by 0.23dB (0.11 standard error).

References

- Steiner, T.J.; Scher, A.I.; Stewart, W.F.; Kolodner, K.; Lieberman, J.; Lipton, R.B. The prevalence and disability burden of adult migraine in England and their relationships to age, gender and ethnicity. *Cephalalgia* **2003**, *23*(7), 519–527.
- Committee, I.H.S.I.H.C. Classification and diagnostic criteria for headache disorders, cranial neuralgias and facial pain. *Cephalalgia* **2013**, *33*, 629–808.
- Schürks, M.; Buringm, J.E.; Kurth, T. Migraine, migraine features and cardiovascular disease. *Headache* **2010**, *50*(6), 1031–1040.
- Lipton, R.B.; Diamond, S.; Reed, M.; Diamond, M.L.; Stewart, W.F. Migraine diagnosis and treatment: Results from the American Migraine Study II. *Headache* **2001**, *41*, 638–645.
- Marcus, D.A.; Soso, M.J. Migraine and stripe-induced visual discomfort. *Archives of Neurology* **1989**, *46*(10), 1129–1132.
- O'Hare, L.; Hibbard, P.B. Visual processing in migraine. *Cephalalgia* **2016**, *36*(11), 1057–1076.

7. Antal, A.; Temme, J.; Nitsche, M.A. Altered motion perception in migraineurs: Evidence for intercortical hyperexcitability. *Cephalalgia* **2005**, *25*, 788–794.
8. McKendrick, A.M.; Badcock, D.R.; Gurgone, M. Vernier acuity is normal in migraine, whereas global form and motion perception are not. *Investigative Ophthalmology and Visual Science* **2006**, *47*, 3213–3219.
9. Ditchfield, J.A.; McKendrick, A.M.; Badcock, D.R. Processing of global form and motion in migraineurs. *Vision Research* **2006**, *46*(1-2), 141–148.
10. Webster, K.E.; Dickinson, J.E.; Battista, J.; McKendrick, A.M.; Badcock, D.R. Increased internal noise cannot account for motion coherence processing deficits in migraine. *Cephalalgia* **2011**, *31*, 1199–1210.
11. Braunitzer, G.; Roksin, A.; Kobor, J.; Benedek, G.; Ngy, A.; Kincses, Z.T. Delayed development of visual motion processing in childhood migraine. *Cephalalgia* **2012**, *32*, 492–496.
12. Shepherd, A.J.; Beaumont, H.M.; Hine, T.J. Motion processing deficits in migraine are related to contrast sensitivity. *Cephalalgia* **2012**, *32*(7), 554–570.
13. Tibber, M.S.; Kelly, M.; Jansari, A.; Dakin, S.C.; Shepherd, A.J. An inability to exclude visual noise in migraine. *Investigative Ophthalmology and Visual Science* **2014**, *55*, 2539–2546.
14. Dakin, S.C.; Mareschal, I.; Bex, P.J. Local and global limitations on direction integration assessed using equivalent noise analysis. *Vision Research* **2005**, *45*, 3027–3049.
15. Geisler, W.S. Motion streaks provide a spatial code for motion direction. *Nature* **1999**, *400*, 65–69.
16. Burr, D.C.; Ross, J. Direct evidence that speedlines influence motion mechanisms. *The Journal of Neuroscience* **2002**, *22*(19), 8661–8664.
17. Athorp, D.; Wenderoth, P.; Alais, D. Motion streaks in fast motion rivalry cause orientation-selective suppression. *Journal of Vision* **2009**, *9*(5), 1–14.
18. Athorp, D.; Cass, J.; Alais, D. Orientation tuning of contrast masking caused by motion streaks. *Journal of Vision* **2010**, *10*(10), 1–13.
19. Athorp, D.; Cass, J.; Alais, D. The spatial tuning of motion streak mechanisms revealed by masking and adaptation. *Journal of Vision* **2011**, *11*(7), 1–16.
20. Alais, D.; Athorp, D.; Karmann, A.; Cass, J. Temporal integration of movement: the time-course of motion streaks revealed by masking. *PLoS ONE* **2011**, *6*(12), e28675.
21. Zhaoping, L. *Understanding Vision: Theory, Models and Data*; Oxford University Press: Oxford, 2014.
22. Series, P.; Latham, P.E.; Pouget, A. Tuning curve sharpening for orientation selectivity: coding efficiency and the impact of correlations. *Nature Neuroscience* **2004**, *7*, 1129–1135.
23. Edwards, M.; Crane, M.F. Motion streaks improve detection. *Vision Research* **2007**, *47*, 828–833.
24. Rogalinski, S.; Rambold, H.A. Probing early motion processing with eye movements: Differences of vestibular migraine, migraine with and without aura in headache-free interval. *Headache* **2017**, *58*(2), 275–286.
25. Drummond, P.D. Motion sickness and migraine: optokinetic stimulation increases scalp tenderness, pain sensitivity in the fingers and photophobia. *Cephalalgia* **2002**, *22*(2), 117–24.
26. Takahashi, M.; Igarashi, M. Comparison of vertical and horizontal optokinetic nystagmus in the squirrel monkey. *ORL* **1977**, *39*, 321–329.
27. Takahashi, M.; Sakurai, S.; Kanzaki, J. Horizontal and vertical optokinetic nystagmus in man. *ORL* **1979**, *40*, 43–52.
28. Lenth, R.V. Least-squares means: The R package lsmeans. *Journal of Statistical Software* **2016**, *69*(1), 1–33.
29. Winter, B. Linear models and linear mixed effects models in R with linguistic applications. *Computation and Language* **2013**, *v1*, arXiv:1308.5499.
30. Calandre, E.P.; Bembibre, J.; Arnedo, M.L.; Becerra, D. Cognitive disturbances and regional cerebral blood flow abnormalities in migraine patients: their relationship with the clinical manifestations of the illness. *Cephalalgia* **2002**, *22*(4), 291–302.
31. Lang, E.; Kaltenhäuser, M.; Neundörfer, B.; Siedler, S. Hyperexcitability of the primary somatosensory cortex in migraine - a magnetoencephalographic study. *Brain* **2004**, *127*, 2459–69.
32. Suhr, J.A.; Seng, E.K. Neurophysiological functioning in migraine: clinical and research implications. *Cephalalgia* **2012**, *32*, 39–54.
33. Salman, A.G.; Hamid, M.A.A.; Mansour, D.E. Correlation of visual field defects and optical coherence tomography finding in migraine patients. *Saudi J Ophthalmol* **2015**, *29*(1), 76–80.

34. Yener, A.U.; Korucu, O. Visual field losses in patients with migraine with and without aura and tension-type headache. *Neuroophthalmology* **2017**, *41*(2), 59–67.
35. Nguyen, B.N.; Vingrys, A.J.; McKendrick, A.M. The effect of duration post-migraine on visual electrophysiological and visual field performance in people with migraine. *Cephalalgia* **2014**, *34*(1), 42–57.
36. Khalil, N.M. Investigations of visual function in migraine using visual evoked potentials and visual psychophysical tests. PhD thesis, University of London, London, 1991.
37. Siniatchkin, M.; Averkina, N.; F., A.; Stephani, U.; Gerber, W.D. Neurophysiological reactivity before a migraine attack. *Neuroscience Letters* **2006**, *400*(1-2), 121–4.
38. Sand, T.; Zhitniy, N.; White, L.R.; Stovner, L.J. Visual evoked potential latency, amplitude and habituation in migraine: a longitudinal study. *Clinical Neurophysiology* **2008**, *119*(5), 1020–7.
39. Sand, T.; White, L.R.; Hagen, K.; Stovner, L.J. Visual evoked potential and spatial frequency in migraine: a longitudinal study. *Acta Neurol Scand Suppl* **2009**, *189*, 33–7.
40. Karovanic, O.; Thabet, M.; Wilson, H.R.; Wilkinson, F. Detection and discrimination of flicker contrast in migraine. *Cephalalgia* **2011**, *31*(6), 723–36.
41. Thabet, M.; Wilkinson, F.; Wilson, H.R.; Karanovic, O. The locus of flicker adaptation in the migraine visual system: A dichoptic study. *Cephalalgia* **2013**, *33*(1), 5–19.
42. Coleston, D.M.; Kennard, C. Responses to temporal visual stimulation in migraine: the critical flicker fusion test. *Cephalalgia* **1995**, *15*(5), 396–8.
43. Kowacs, P.A.; Piovesan, E.J.; Werneck, L.C.; Famelie, H.; Zani, A.N.; da Silva, H.P. Critical flicker frequency in migraine: A controlled study in patients without prophylactic therapy. *Cephalalgia* **2005**, *25*(5), 339–43.
44. Shepherd, A.J.; Wyatt, G.; Tibber, M.S. Visual metacontrast masking in migraine. *Cephalalgia* **2011**, *31*(3), 346–356.
45. Schoenen, J.; Wang, W.; Albert, A.; Delwaide, P.J. Potentiation instead of habituation characterizes visual evoked potentials in migraine patients between attacks. *European Journal of Neuroscience* **1995**, *2*, 115–22.
46. Hodkinson, D.J.; Veggeberg, R.; Kucyi, A.; van Dijk, K.R.A.; Wilcox, S.L.; Scrivani, S.J.; Burstein, R.; Becerra, L.; Borsook, D. Cortico-cortical connections of primary sensory areas and associated symptoms in migraine. *eNeuro* **2016**, *3*(6).
47. Tedeschi, G.; Russo, A.; Conte, F.; Caiazzo, G.; Giordano, A.; Conforti, R.; Esposito, F.; Tessitore, A. Increased interictal visual network connectivity in patients with migraine with visual aura. *Cephalalgia* **2016**, *36*(2), 139–47.
48. Keil, J.; Müller, N.; Ihssen, N.; Weisz, N. On the variability of the McGurk effect: audiovisual integration depends on prestimulus brain states. *Cerebral Cortex* **2012**, *22*(1), 221–31.
49. Kumar, V.G.; Halder, T.; Jaiswal, A.K.; Mukherjee, A.; Roy, D.; Banerjee, A. Large scale functional brain networks underlying temporal integration of audio-visual speech perception: An EEG study. *Frontiers in Psychology* **2016**, *7*, 1558.
50. de Tommaso, M.; Stamaglia, S.; Marinazzo, D.; Trotta, G.; Pellicoro, M. Functional and effective connectivity in EEG alpha and beta bands during intermittent flash stimulation in migraine with and without aura. *Cephalalgia* **2013**, *33*, 938–947.
51. Jensen, O.; Gips, B.; Bergmann, T.O.; Bonneford, M. Temporal coding organized by coupled alpha and gamma oscillations prioritize visual processing. *Trends in Neurosciences* **2014**, *37*(7), 357–369.
52. Dugue, L.; MARque, P.; VanRullen, R. The phase of ongoing oscillations mediates the causal relation between brain excitation and visual perception. *Journal of Neuroscience* **2011**, *31*(33), 11893–11889.
53. Ergenoglu, T.; Demiralp, T.; Bayrataroglu, Z.; Ergen, M.; Beydagi, H.; Uresin, Y. Alpha rhythm of the EEG modulates visual detection performance in humans. *Cognitive Brain Research* **2004**, *20*, 376–383.
54. Hanslmayr, S.; Klimesch, W.; Sauseng, P.; Gruber, W.; Doppelmayr, M.; Freunberger, R.; Pecherstorfer, T. Visual discrimination performance is related to decreased alpha amplitude but increased phase locking. *Neuroscience Letters* **2005**, *375*, 64–68.
55. Hanslmayr, S.; Aslan, A.; Staudigl, T.; Klimesch, W.; Herrmann, C.S.; Bauml, K.H. Prestimulus oscillations predict visual perception performance between and within subjects. *NeuroImage* **2007**, *37*, 1465–1473.
56. Cecere, R.; Rees, G.; Romei, V. Individual differences in alpha frequency drive crossmodal illusory perception. *Current Biology* **2015**, *25*, 231–235.
57. Zaehle, T.; Rach, S.; Herrmann, C.S. Transcranial alternating current stimulation enhances alpha activity in human EEG. *PLoS ONE* **2010**, *5*(11), e13766.

58. Yang, W.; Chu, B.; Yang, J.; Yu, Y.; Wu, J.; Yu, S. Elevated audiotemporal interaction in patients with migraine without aura. *The Journal of Headache and Pain* **2014**, *15*, 44.
59. de Tommaso, M.; Ambrosini, A.; Brighina, F.; Coppola, G.; Perrotta, A.; Pierelli, F.; Sandrini, G.; Valeriani, M.; Marinazzo, D.; Stramaglia, S.; Schoenen, J. Altered processing of sensory stimuli in patients with migraine. *Nature Reviews Neuroscience* **2014**, *10*(3), 144–55.
60. Battista, J.; Badcock, D.R.; McKendrick, A.M. Migraine increases centre-surround suppression for drifting visual stimuli. *PLoS ONE* **2011**, *6*(4), e18211.
61. Coppola, G.; Parisi, V.; Lorenzo, C.D.; Serrao, M.; Magis, D.; Schoenen, J. Lateral inhibition in visual cortex of migraine patients between attacks. *Headache Pain* **2013**, *14*(1), 20.
62. Asher, J.M.; O'Hare, L.; Romei, V.; Hibbard, P.B. Typical lateral interactions, but increased contrast sensitivity, in migraine-with-aura. *Vision* **2018**, *2*(1), 7.
63. Adelson, E.H.; Bergen, J.R. Spatiotemporal energy models for the perception of motion. *J Opt Soc Am A* **1985**, *2*(2), 284–299.
64. Granziera, C.; DaSilva, A.F.M.; Snyder, J.; Tuch, D.S.; Hadjikhani, N. Anatomical alterations of the visual motion processing network in migraine with and without aura. *PLoS Med* **2006**, *3*(10), e402.
65. Edmeads, J.; Findlay, H.; Tugwell, P.; Pryse-Phillips, W.; Nelson, R.F.; Murray, T.J. A Canadian population survey on the clinical, epidemiologic and societal impact of migraine and tension-type headache. *Can J Neurol Sci* **1993**, *19*, 333–339.
66. MacGregor, E.A.; Brandes, J.; Eikermann, A. Migraine prevalence and treatment patterns: The Global Migraine and Zolmitriptan Evaluation Survey. *Headache* **2003**, *43*(1), 19–26.
67. McKendrick, A.M.; Badcock, D.R. Motion processing deficits in migraine. *Cephalalgia* **2004**, *24*, 363–372.
68. Battista, J.; Badcock, D.R.; McKendrick, A.M. Centre-surround visual motion processing in migraine. *Invest Ophthalmol Vis Sci* **2010**, *51*, 6070–6076.
69. Coppola, G.; Renzo, A.D.; Tnelli, E.; Lepre, C.; Lorenzo, C.D.; Lorenzo, G.D.; Scapeccia, M.; Parisi, V.; Serrao, M.; Colonnese, C.; Schoenen, J.; Pierelli, F. Thalamo-cortical network activity between migraine attacks: Insights from MRI-based microstructural and functional resting-state network correlation analysis. *Journal of Headache and Pain* **2016**, *17*(1), 100.
70. Coppola, G.; Renzo, A.D.; Tinelli, E.; Lorenzo, C.D.; Lorenzo, G.D.; Parisi, V.; Serrao, M.; Schoenen, J.; Pierelli, F. Thalamo-cortical network activity during spontaneous migraine attacks. *Neurology* **2016**, *87*(20), 2154–2160.
71. Faragó, P.; Tuka, B.; Tóth, E.; Szabó, N.; Király, A.; Csete, G.; Szok, D.; Tajti, J.; Párdutz, A.; Vécsei, L.; Kincses, Z. Interictal brain activity differs in migraine with and without aura: resting state fMRI study. *Journal of Headache and Pain* **2017**, *18*(1), 8.
72. Brainard, D.H. The psychophysics toolbox. *Spatial Vision* **1997**, *10*(4), 433–436.
73. Pelli, D.G. The videotoolbox for visual psychophysics: Transforming numbers into movies. *Spatial Vision* **1997**, *10*(4), 437–442.
74. Kleiner, M.; Brainard, D.; Pelli, D. What's new in Psychtoolbox-3? *Perception* **2007**, *365* (ECVP supplement), S14.
75. R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2013.
76. Bolker, H.S.; Westfall, J.; Aust, F.; R Core Team. *Afex: Analysis of factorial experiments.*, 2017. R package version 0.18-0.
77. Bates, D.; Maechler, M.; bolker, B.; Walker, S. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **2015**, *67*(1), 1–48.
78. Norman, G. Likert scales, levels of measurement and the "laws" of statistics. *Adv Health Sci Edu Theory Pract* **2010**, *15*(5), 625–32.
79. Nimon, K.F. Statistical assumptions of substantive analyses across the general linear model: A mini-review. *Frontiers in Psychology* **2012**, *3*, 322.
80. Khan, A.; Rayner, G.D. Robustness to non-normality of common tests for the many-sample location problem. *Journal of Applied Mathematics and Decision Sciences* **2003**, *7*(4), 187–206.
81. Noguchi, K.; Gel, Y.R.; Brunner, E.; Konietzke, F. nparLD: An R Software Package for the Nonparametric Analysis of Longitudinal Data in Factorial Experiments. *Journal of Statistical Software* **2012**, *50*, 1–23.

- 631 82. Naito, T.; Sato, H.; Osaka, N. Direction anisotropy of human motion perception depends on stimulus
632 speed. *Vision Research* **2010**, *50*(18), 1862–1866.
- 633 83. Ohtani, Y.; Ejima, Y. Anisotropy for direction discrimination in a two-frame apparent motion display.
634 *Vision Research* **1997**, *37*(6), 765–767.

635 © 2018 by the author. Submitted to *Vision* for possible open access publication under the terms and conditions
636 of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).